

Crit Care. 2023;27(1):199. doi:10.1186/s13054-023-04491-w.

15. Mascarenhas M, Beattie M, Roxburgh M, MacKintosh J, Clarke N, Srivastava D. Using the Model for Improvement to implement the Critical-Care Pain Observation Tool in an adult intensive care unit. *BMJ Open Qual.* 2018;7(4):e000304. doi:10.1136/bmjopen-2017-000304.
16. Havaladar AA. Pain Assessment in Intensive Care Unit: A Forgotten Entity or a Quality Indicator? *Indian Journal of Critical Care Medicine.* 2022;26(4):421-422. doi:10.5005/jp-journals-10071-24191.
17. McGovern C, Cowan R, Appleton R, Miles B. Pain, agitation and delirium in the intensive care unit. *Anaesthesia & Intensive Care Medicine.* 2021;22(12):799-806. doi:10.1016/j.mpaic.2021.10.013.
18. Morris JL, Bernard F, Bérubé M, et al. Determinants of pain assessment documentation in intensive care units. *Can J Anesth/J Can Anesth.* 2021;68(8):1176-1184. doi:10.1007/s12630-021-02022-1.
19. Gomarverdi S, Sedighie L, Seifrabiei M, Nikooseresht M. Comparison of two pain scales: Behavioral pain scale and critical-care pain observation tool during invasive and noninvasive procedures in intensive care unit-admitted patients. *Iranian J Nursing Midwifery Res.* 2019;24(2):151. doi:10.4103/ijnmr.IJNMR_47_18.
20. Hora TCNSD, Alves IGN. Scales for the assessment of pain in the intensive care unit. Systematic review. *Brazilian Journal Of Pain.* 2020;3(3). doi:10.5935/2595-0118.20200043.
21. Erden S, Demir N, Ugras GA, Arslan U, Arslan S. Vital signs: Valid indicators to assess pain in intensive care unit patients? An observational, descriptive study. *Nursing & Health Sciences.* 2018;20(4):502-508. doi:10.1111/nhs.12543.
22. Severgnini P, Pelosi P, Contino E, Serafinelli E, Novario R, Chiaranda M. Accuracy of Critical Care Pain Observation Tool and Behavioral Pain Scale to assess pain in critically ill conscious and unconscious patients: prospective, observational study. *J intensive care.* 2016;4(1):68. doi:10.1186/s40560-016-0192-x.
23. Chamorro C. Pain in the ICU – The fifth sign, not the fifth element. *Medicina Intensiva.* 2016;40(8):461-462. doi:10.1016/j.medin.2016.08.010.
24. Rijkenberg S, van der Voort PHJ. Can the critical-care pain observation tool (CPOT) be used to assess pain in delirious ICU patients? *J Thorac, Dis.* 2016;8(5):E285-E287. doi:10.21037/jtd.2016.03.32.
25. Álvarez EA, Parada FJ. Association of Pain During the Evaluation of Delirium in Intensive Care Unit Patients. *Front Med.* 2021;8:722001. doi:10.3389/fmed.2021.722001.
26. Fischer T, Hosie A, Luckett T, Agar M, Phillips J. Strategies for Pain Assessment in Adult Patients With Delirium: A Scoping Review. *Journal of Pain and Symptom Management.* 2019;58(3):487-502.e11. doi:10.1016/j.jpainsymman.2019.05.020.
27. Sampson EL, West E, Fischer T. Pain and delirium: mechanisms, assessment, and management. *Eur Geriatr Med.* 2020;11(1):45-52. doi:10.1007/s41999-019-00281-2.
28. Sadaf F, Saqib M, Iftikhar M, Ahmad A. Prevalence and Risk Factors of Delirium in Patients Admitted to Intensive Care Units: A Multicentric Cross-Sectional Study. *Cureus.* Published online September 7, 2023. doi:10.7759/cureus.44827.
29. Renzi S, Gitti N, Piva S. Delirium in the intensive care unit: a narrative review. *JGG.* 2023;71(1):22-29. doi:10.36150/2499-6564-N600.
30. La Cour KN, Andersen-Ranberg NC, Weihe S, et al. Distribution of delirium motor subtypes in the intensive care unit: a systematic scoping review. *Crit Care.* 2022;26(1):53. doi:10.1186/s13054-022-03931-3.
31. Alaterre C, Fazilleau C, Cayot-Constantin S, et al. Monitoring delirium in the intensive care unit: Diagnostic accuracy of the CAM-ICU tool when performed by certified nursing assistants – A prospective multicenter study. *Intensive and Critical Care Nursing.* 2023;79:103487. doi:10.1016/j.iccn.2023.103487.
32. Stollings JL, Kotfis K, Chanques G, Pun BT, Pandharipande PP, Ely EW. Delirium in critical illness: clinical manifestations, outcomes, and management. *Intensive Care Med.* 2021;47(10):1089-1103. doi:10.1007/s00134-021-06503-1.
33. Gao Y, Gong S, Zhou W, Li X, Gan X. Frequency and Risk Factors of Subsyndromal Delirium in the Intensive Care Units: A Prospective Cohort Study. *NDT.* 2023;Volume 19:1003-1016. doi:10.2147/NDT.S407156.
34. Mart MF, Williams Roberson S, Salas B, Pandharipande PP, Ely EW. Prevention and Management of Delirium in the Intensive Care Unit. *Semin Respir Crit Care Med.* 2021;42(01):112-126. doi:10.1055/s-0040-1710572.
35. Lobo-Valbuena B, Gordo F, Abella A, et al. Risk factors associated with the development of delirium in general ICU patients. A prospective observational study. *Zivkovic AR, ed. PLoS ONE.* 2021;16(9):e0255522. doi:10.1371/journal.pone.0255522.
36. Ormseth CH, LaHue SC, Oldham MA, Josephson SA, Whitaker E, Douglas VC. Predisposing and Precipitating Factors Associated With Delirium: A Systematic Review. *JAMA Netw Open.* 2023;6(1):e2249950. doi:10.1001/jamanetworkopen.2022.49950.
37. Gagnon PR. Treatment of delirium in supportive and palliative care: Current Opinion in Supportive and Palliative Care. 2008;2(1):60-66. doi:10.1097/SPC.0b013e3282f4ce05.
38. Bouajram RH, Sebat CM, Love D, Louie EL, Wilson MD, Duby JJ. Comparison of Self-Reported and Behavioral Pain Assessment Tools in Critically Ill Patients. *J Intensive Care Med.* 2020;35(5):453-460. doi:10.1177/0885066618757450.
39. Sun G, Wen Z, Ok D, Doan L, Wang J, Chen ZS. Detecting acute pain signals from human EEG. *Journal of Neuroscience Methods.* 2021;347:108964. doi:10.1016/j.jneumeth.2020.108964.
40. Modares-Haghighi P, Boostani R, Nami M, Sanei S. Quantification of pain severity using EEG-based functional connectivity. *Biomedical Signal Processing and Control.* 2021;69:102840. doi:10.1016/j.bspc.2021.102840.
41. Zis P, Liampas A, Artemiadis A, et al. EEG Recordings as Biomarkers of Pain Perception: Where Do We Stand and Where to Go? *Pain Ther.* 2022;11(2):369-380. doi:10.1007/s40122-022-00372-2.
42. Coleman RM, Tousignant-Laflamme Y, Ouellet P, Parenteau-Goudreault É, Cogan J, Bourgault P. The Use of the Bispectral Index in the Detection of Pain in Mechanically Ventilated Adults in the Intensive Care Unit: A Review of the Literature. *Pain Research and Management.* 2015;20(1):e33-e37. doi:10.1155/2015/981419.
43. Tasaka CL, Duby JJ, Pandya K, Wilson MD, A. Hardin K. Inadequate Sedation During Therapeutic Paralysis: Use of Bispectral Index in Critically Ill Patients. *Drugs - Real World Outcomes.* 2016;3(2):201-208. doi:10.1007/s40801-016-0076-3.
44. Faritous Z, Barzanji A, Azarfarin R, et al. Comparison of Bispectral Index Monitoring With the Critical-Care Pain Observation Tool in the Pain Assessment of Intubated Adult Patients After Cardiac Surgery. *Anesth Pain Med.* 2016;6(4). doi:10.5812/aapm.38334.
45. Bower MM, Sweidan AJ, Xu JC, Stern-Nezer S, Yu W, Groysman LI. Quantitative Pupillometry in the Intensive Care Unit. *J Intensive Care Med.* 2021;36(4):383-391. doi:10.1177/0885066619881124.
46. Arican Ş. Comparison of pupillometer, critical care pain observation scale and vital findings in the evaluation of pain in intensive care patients without oral communication. *Agri.* Published online 2022. doi:10.14744/agri.2021.72246.
47. Singh A. A, Kumar N, De RR, Bahadur R, Shekhar S. The Role of Pupillometry in the Assessment of Pain in Children Under General Anesthesia: A Prospective Single-Blinded Observational Study. *Cureus.* Published online August 22, 2023. doi:10.7759/cureus.43894.
48. Aissou M, Snauwaert A, Dupuis C, Atchabahan A, Aubrun F, Beaussier M. Objective Assessment of the Immediate Postoperative Analgesia Using Pupillary Reflex Measurement. *Anesthesiology.* 2012;116(5):1006-1012. doi:10.1097/ALN.0b013e318251d1fb.
49. Frattino S, Peluso L, Talamonti M, et al. Evaluation of Nociception Using Quantitative Pupillometry and Skin Conductance in Critically Ill Unconscious Patients: A Pilot Study. *Brain Sciences.* 2021;11(1):109. doi:10.3390/brainsci11010109.
50. Brett K, Severn M. Facial Analysis Technology for Pain Detection: A Potentially Useful Tool for People Living With Dementia. *cjht.* 2023;3(7). doi:10.51731/cjht.2023.682.
51. Pu L, Coppieters MW, Smalbrugge M, et al. Implementing PainChek and PARO to Support Pain Assessment and Management in Residents with Dementia: A Qualitative Study. *Pain Management Nursing.* 2023;24(6):587-594. doi:10.1016/j.pmn.2023.04.001.
52. Crellin DJ, Harrison D, Santamaria N, Babl FE. Comparison of the Psychometric Properties of the FLACC Scale, the MBPS and the Observer Applied Visual Analogue Scale Used to Assess Procedural Pain. *JPR.* 2021;Volume 14:881-892. doi:10.2147/JPR.S267839.
53. Lee S, Jung DE, Park D, et al. Intraoperative neurological pupil index and postoperative delirium and neurologic adverse events after cardiac surgery: an observational study. *Sci Rep.* 2023;13(1):13838. doi:10.1038/s41598-023-41151-z.
54. Mitra S, Saxena P, Jain K, et al. Clinical Utility of the Behavioral Pain Assessment Tool in Patients Admitted in the Intensive Care Unit. *Indian Journal of Critical Care Medicine.* 2020;24(8):695-700. doi:10.5005/jp-journals-10071-23521.
55. Nugent SM, Lovejoy TI, Shull S, Dobscha SK, Morasco BJ. Associations of Pain Numeric Rating Scale Scores Collected during Usual Care with Research Administered Patient Reported Pain Outcomes. *Pain Medicine.* 2021;22(10):2235-2241. doi:10.1093/pm/pnab110.
56. Wiegand DL, Wilson T, Pannullo D, et al. Measuring Acute Pain Over Time in the Critically Ill Using the Multidimensional Objective Pain Assessment Tool (MOPAT). *Pain Management Nursing.* 2018;19(3):277-287. doi:10.1016/j.pmn.2017.10.013.